



Food Microbiology

Production of docosahexaenoic acid by *Aurantiochytrium* sp. ATCC PRA-276

Valcenir Júnior Mendes Furlan^{a,*}, Victor Maus^b, Irineu Batista^c,
Narcisa Maria Bandarra^c

^a Universidade Federal do Pampa (UNIPAMPA), Itaqui, RS, Brazil

^b International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

^c Portuguese Institute of Sea and Atmosphere (IPMA, I.P./DMRM), Lisbon, Portugal

ARTICLE INFO

Article history:

Received 3 August 2015

Accepted 13 October 2016

Available online 20 January 2017

Associate Editor: Rosane Freitas

Schwan

Keywords:

Carbon source

Docosahexaenoic acid

Nitrogen source

Polyunsaturated fatty acids

Thraustochytrids

ABSTRACT

The high costs and environmental concerns associated with using marine resources as sources of oils rich in polyunsaturated fatty acids have prompted searches for alternative sources of such oils. Some microorganisms, among them members of the genus *Aurantiochytrium*, can synthesize large amounts of these biocompounds. However, various parameters that affect the polyunsaturated fatty acids production of these organisms, such as the carbon and nitrogen sources supplied during their cultivation, require further elucidation. The objective of this investigation was to study the effect of different concentrations of carbon and total nitrogen on the production of polyunsaturated fatty acids, particularly docosahexaenoic acid, by *Aurantiochytrium* sp. ATCC PRA-276. We performed batch system experiments using an initial glucose concentration of 30 g/L and three different concentrations of total nitrogen, including 3.0, 0.44, and 0.22 g/L, and fed-batch system experiments in which 0.14 g/L of glucose and 0.0014 g/L of total nitrogen were supplied hourly. To assess the effects of these different treatments, we determined the biomass, glucose, total nitrogen and polyunsaturated fatty acids concentration. The maximum cell concentration (23.9 g/L) was obtained after 96 h of cultivation in the batch system using initial concentrations of 0.22 g/L total nitrogen and 30 g/L glucose. Under these conditions, we observed the highest level of polyunsaturated fatty acids production (3.6 g/L), with docosahexaenoic acid and docosapentaenoic acid ω_6 concentrations reaching 2.54 and 0.80 g/L, respectively.

© 2017 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

The search for nutraceutical products that can prevent and/or treat diseases has intensified during the last decade.

Among these products, types ω_3 and ω_6 polyunsaturated fatty acids (PUFAs) have received a great deal of attention due to their health benefits and their extensive applications in the food and pharmaceutical industries.¹ Docosahexaenoic acid (DHA, C22:6 ω_3), for example, is necessary for the brain development of newborn children and contributes to increasing their intelligence and verbal and reasoning skills.² Furthermore, DHA is helpful in treating

* Corresponding author.

E-mail: juniorfurlan@yahoo.com.br (V.J. Furlan).

<http://dx.doi.org/10.1016/j.bjm.2017.01.001>

1517-8382/© 2017 Sociedade Brasileira de Microbiologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

atherosclerosis, rheumatoid arthritis, and Alzheimer's disease,³ as well as in preventing breast and colon cancer.⁴ Docosapentaenoic acid (DPA, C22:5 ω6) is another PUFA that is important for human health. DPA has been found to help prevent various diseases, such as cardiovascular disorders (such as myocardial infarction, thrombosis, and atherosclerosis), diabetes, asthma, inflammation and rheumatism (including arthritis and osteoporosis).⁵

The main commercial source of these compounds, particularly DHA, is oil obtained from marine fish.⁶ However, the widespread consumption of these oils is limited by marine chemical pollution, declining fish stocks, seasonal variations in the composition of fish oils, their poor oxidative stability, typical unpleasant odour and taste, and the high cost of their extraction and purification processes.⁷ These problems have inspired the development of new methods for the large-scale production of oils by safe and healthy sources.^{1,8}

Heterotrophic microorganisms that are members of the Thraustochytrid group are alternative sources of these oils. These oleaginous microorganisms can accumulate more than 50% of their weight as lipids, with a high concentration of DHA of greater than 25% of the total lipids.⁹ Furthermore, the lipids of thraustochytrids contain a specific PUFA (DHA) instead of a mixture of PUFAs. Therefore, their oil has a higher level of oxidative stability than that of fish oil.¹⁰ The appropriate concentrations of carbon and nitrogen are essential for thraustochytrids to biosynthesize and accumulate polyunsaturated fatty acids. The concentration of the carbon source affects the synthesis of organic molecules and the availability of energy, whereas the concentration of the nitrogen source affects the synthesis of amino acids and nucleic acids. Therefore, understanding the effects of these substrates on cultivated microorganisms is crucial for optimizing their oil production. Herein, we present the results of a study of the effect of different concentrations of carbon and nitrogen on the PUFAs production of *Aurantiochytrium* sp. ATCC PRA-276.

Materials and methods

Microorganism

Aurantiochytrium sp. ATCC PRA-276 cells were obtained from the American Type Culture Collection (Manassas, VA, USA).

Preparing the inoculum

Aurantiochytrium sp. ATCC PRA-276 cells grown on potato dextrose agar and stored at 4 °C were transferred to 500 mL flasks containing 100 mL of culture medium consisting (g/L) of yeast extract (1.0), peptone (15.0) and glucose (20.0) dissolved in seawater (1.5% w/v). The glucose stock solution was sterilized separately. The cells were incubated in an orbital shaker (Ika, KS 260B) rotating at 150 rpm at 30 °C without light for 48 h.¹¹

Preparing the culture media

We cultivated the microorganism in a Biostat® Bplus bioreactor (Sartorius Stedim Biotech., Germany) containing a

5L borosilicate glass vessel and equipped with pressure flow meters and gas and liquid-flow controllers. We conducted batch and fed-batch experiments. For the batch system experiments we used a medium consisting of KH₂PO₄ (0.3 g/L), MgSO₄·7H₂O (5.0 g/L), NaCl (10.0 g/L), NaHCO₃ (0.1 g/L), CaCl₂·2H₂O (0.3 g/L), KCl (0.28 g/L), glucose (30.0 g/L) and different total nitrogen (TN) concentrations: 3.0 g/L (1.36 g/L (NH₄)₂SO₄, 13.63 g/L yeast extract and 13.63 g/L monosodium glutamate), 0.44 g/L (0.2 g/L (NH₄)₂SO₄, 2.0 g/L yeast extract and 2.0 g/L monosodium glutamate), and 0.22 g/L (0.1 g/L de (NH₄)₂SO₄, 1.0 g/L yeast extract and 1.0 g/L monosodium glutamate). For the fed-batch system experiments, 0.14 g/L of glucose and 0.0014 g/L of total nitrogen were supplied hourly (6.66 × 10⁻⁴ g/L h of (NH₄)₂SO₄, 6.66 × 10⁻³ g/L h of yeast extract and 6.66 × 10⁻³ g/L h of monosodium glutamate). The yeast extract, monosodium glutamate and glucose solutions were separately sterilized by treatment at 121 °C for 15 min in a CerroClav CV-EL-18 L autoclave. The bioreactor was sterilized in an AJC Uniclave 77-127 L autoclave for 60 min. The other components of the medium were filtered-sterilized using 0.22 μm membranes (Millipore).

After sterilization, the dissolved components were added to the bioreactor along with the following metal solutions: MnCl₂·4H₂O (8.6 mg/L), ZnCl₂ (0.6 mg/L), CoCl₂·4H₂O (0.26 mg/L), CuSO₄·5H₂O (0.02 mg/L), FeCl₃·6H₂O (2.9 mg/L), H₃BO₃ (34.2 mg/L), and Na₂EDTA (30.0 mg/L) and the following vitamin solutions: thiamine (9.5 mg/L) and calcium pantothenate (3.2 mg/L), all of which had been sterilized using 0.22 μm membrane filters (Millipore). The inoculum (350 mL) was then added to the culture medium (10% v/v relative to the total volume of the culture medium).

Cultivation was performed at 23 °C with agitation at 100 rpm and the pH of the media adjusted to 6.0 using 4N NaOH. During the first 120 h of cultivation, the culture medium was aerated at 2.5 vvm. After this period, air injection was discontinued.

Biomass concentration

The biomass was determined every 24 h using the method of Min et al.¹² An aliquot of the culture medium was filtered using a previously weighed glass-microfiber filter paper (GF/C: 1.2 μm, Whatman). The biomass retained in the filter was washed twice using distilled water and dried in an oven (Memmert) at 60 °C for 24 h. The biomass content was calculated as the difference between the initial and final weights.

Glucose concentration

The glucose content of the culture supernatant was determined each 24 h using the spectrophotometric method described by Miller,¹³ using a UV/Vis dual beam absorption spectrophotometer (Ati Unicam Helios Alpha, UK).

Total nitrogen concentration

The total nitrogen (defined and complex sources) content of the culture supernatant was determined each 24 h following the procedure described by Furlan et al.¹¹

Fatty acid profile

Culture samples collected at 24 h intervals were centrifuged (Kubota, 6800) at $8742 \times g$ for 15 min at 4°C , after which the biomass was washed with distilled water and centrifuged again. This process was repeated twice. The biomass was frozen at -20°C and dried for 48 h using a freeze dryer (Heto PowerDry LL 3000).

Between 20 and 100 mg of the lyophilized biomass was weighed and added to $50 \mu\text{L}$ of the internal C21:0 standard solution (10 mg/mL) to permit expressing the results as g of fatty acids/g of lyophilized biomass.

Methyl esters of the fatty acids were prepared by esterification using the acid catalysis method described by Cohen et al.¹⁴ A gas chromatography system (Varian, CP 3800) equipped with an autosampler, injector, and flame ionization detector (FID), both of the latter at 250°C , was used to identify the methyl esters in the samples. The methyl esters were separated using a DB-WAX polyethylene glycol capillary column (Agilent, 30 m long, 0.25 mm internal diameter and 0.25 μm thick) using the following program: heating at 180°C (5 min), gradually increasing at $4^\circ\text{C}/\text{min}$ to 220°C (and holding for 25 min), and then gradually increasing at $20^\circ\text{C}/\text{min}$ to 240°C (and holding for 15 min). The methyl esters were identified by comparing their retention times with those of chromatographic standards (Sigma-Aldrich Co, St. Louis, MO, USA).

The results were analyzed using an analysis of variance (ANOVA) and the mean values were compared using the Tukey test, with the significance level set at 5%. Before performing the ANOVA, the normality of the data distributions

were evaluated using the Kolmogorov–Smirnov test and the homoscedasticity of the data was evaluated using the Cochran test.¹⁵

Results

Growth kinetics, glucose and total nitrogen consumption

Fig. 1 shows the average biomass, glucose and total nitrogen concentrations over time in cultures of *Aurantiochytrium* sp. ATCC PRA-276 growing under four different conditions.

As shown in Fig. 1(A), in the experiments using an initial total nitrogen concentration of 3.0 g/L , the maximum biomass concentration (9.3 g/L) was reached at 120 h, for an average yield of 0.07 g/Lh of biomass. The average glucose consumption rate in these experiments was 0.13 g/Lh and 0.43 g of biomass was produced per gram of glucose consumed ($Y_{\text{Biomass}/\text{Glucose}}$). The average total nitrogen consumption rate was 0.0010 g/Lh , resulting in a substrate to biomass conversion factor ($Y_{\text{Biomass}/\text{nitrogen}}$) of 65.8.

In the experiments using an initial nitrogen concentration of 0.44 g/L , the highest biomass concentration (17.0 g/L) was obtained at 144 h, for an average yield of 0.11 g/Lh of biomass (Fig. 1(B)). The average glucose consumption rate was 0.15 g/Lh , resulting in a glucose to biomass conversion factor ($Y_{\text{Biomass}/\text{glucose}}$) of 6.5. The average total nitrogen consumption rate was 0.0018 g/Lh . Each gram of nitrogen that was consumed was converted into 50.7 g of biomass ($Y_{\text{Biomass}/\text{Nitrogen}}$).




Fig. 1 – Concentrations of biomass, glucose and total nitrogen over time in the culture media of *Aurantiochytrium* sp. ATCC PRA-276. The graphs show the data obtained using different treatments, as follows: a batch system with an (A) initial nitrogen concentration of 3.0 g/L , (B) initial nitrogen concentration of 0.44 g/L or (C) initial nitrogen concentration of 0.22 g/L or (D) a fed-batch system with 0.14 g/L of glucose and 0.0014 g/L of nitrogen supplied hourly.




Fig. 2 – PUFAs concentrations in the biomass of *Aurantiochytrium* sp. ATCC PRA-276 cultivated under different conditions. PUFAs, polyunsaturated fatty acids; TN, total nitrogen.

In the experiments using an initial nitrogen concentration of 0.22 g/L, the highest biomass concentration (23.9 g/L) was observed at 96 h (Fig. 1(C)), for a maximum yield of 0.23 g/L h of biomass. In these experiments, the average glucose consumption rate was 0.17 g/L h and the glucose to biomass conversion factor ($Y_{\text{Biomass/Glucose}}$) was 1.28. The average nitrogen consumption rate was 0.0022 g/L h, resulting in a $Y_{\text{Biomass/Nitrogen}}$ conversion factor of 104.7, meaning that 104.7 g of biomass was produced per gram of nitrogen consumed.

When the fed-batch cultivation process was used, the highest biomass concentration (13.2 g/L) was reached at 120 h (Fig. 1(D)), for a maximum yield of 0.10 g/L h of biomass. The average glucose consumption rate was 0.13 g/L h, resulting in a glucose to biomass conversion factor ($Y_{\text{Biomass/Glucose}}$) of 0.64. Using this cultivation process, the average total nitrogen consumption rate was 0.0016 g/L h and the $Y_{\text{Biomass/Nitrogen}}$ conversion factor was 49.15, meaning that 49.15 g of biomass was produced per gram of nitrogen consumed.

Fatty acid profile

Fig. 2 shows the average PUFAs concentrations in the biomass that were reached throughout *Aurantiochytrium* sp. ATCC PRA-276 cultivation in the different experiments.

Evaluating the PUFAs concentrations in the biomass (g/L) showed that there were significant differences among the values at different times throughout cultivation in all of the experiments. The highest PUFAs concentration (3.6 g/L) was

observed at 96 h of cultivation in the experiments using an initial nitrogen concentration of 0.22 g/L. The second highest PUFAs concentration (2.85 g/L) was obtained at 168 h of cultivation in the experiments using an initial nitrogen concentration of 0.44 g/L. In the fed-batch culture, the maximal PUFAs concentration of 1.89 g/L was reached at 144 h of cultivation. In the experiments using an initial nitrogen concentration of 3 g/L, the maximal PUFAs concentration of 0.84 g/L was reached at 120 h of cultivation (Fig. 2).

Fig. 3 shows the fatty acid profile at the time point when the highest PUFAs yield was obtained in each experiment (g/L).

In the experiments using an initial nitrogen concentration of 3.0 g/L, 9% (w/w) of the biomass was composed of PUFAs at 120 h of cultivation, of which 20% was DPA ω 6 (Fig. 3), which represented 1.8% of the biomass (0.17 g/L). As shown in Fig. 3, DHA accounted for 61.3% of the PUFAs present, i.e., 5.5% of the biomass (0.51 g/L).

Using an initial nitrogen concentration of 0.44 g/L, 18% (w/w) of the biomass was composed of PUFAs at 168 h of cultivation, of which 22% was DPA ω 6, i.e., 3.9% of the biomass (0.63 g/L). We also observed that DHA accounted for 69.2% of the PUFAs, which was 12.5% of the total biomass (1.97 g/L) (Fig. 3).

In the experiments using an initial nitrogen concentration of 0.22 g/L, 15% (w/w) of the biomass was composed of PUFAs at 96 h of cultivation, of which 22.3% was DPA ω 6 (Fig. 3), accounting for 3.34% of the total biomass (0.80 g/L). We also observed




Fig. 3 – Fatty acid profiles of the biomass of *Aurantiochytrium* sp. ATCC PRA-276 when batch-cultured for 120 h using an initial nitrogen concentration of 3.0 g/L total nitrogen, at 168 h when batch-cultured using an initial nitrogen concentration of 0.44 g/L, at 96 h when batch-cultured using an initial nitrogen concentration of 0.22 g/L, and at 144 h when cultured using the fed-batch process.

that DHA comprised 70.5% of the PUFAs (Fig. 3), i.e., 10.62% of the biomass (2.54 g/L).

After 144 h of cultivation using the fed-batch system, 14.9% (w/w) of the biomass was composed of PUFAs, of which 23.1% was DPA ω 6 (Fig. 3), i.e., 3.43% of the total biomass (0.44 g/L). In addition, DHA accounted for 70.5% of the PUFAs (Fig. 3), i.e., 10.5% of the total biomass (1.33 g/L).

Discussion

In the experiments using an initial nitrogen concentration of 3 g/L, the cells consumed the least amount of TN (0.0010 g/L h), resulting in the lowest cell yield (0.07 g/L h). This result could be due to the low C/N substrate ratio in the culture (4). An excess of nitrogen may have inhibited the growth of the evaluated strain. Chen et al.¹⁶ conducted a study in which they optimized the nitrogen sources for *Aurantiochytrium* sp. BR-MP4-A1 and obtained a maximum biomass concentration (9.27 g/L) by supplying 2.4 g/L of TN (as monosodium glutamate, yeast extract, and tryptone), which resulted in a C/N ratio (5). In the experiments in which we used an initial nitrogen concentration of 3 g/L, a similar maximum biomass concentration of 9.3 g/L was reached (Fig. 1(A)).

Using an initial TN concentration of 0.44 g/L to cultivate *Aurantiochytrium* sp. BR-MP4-A1, Li et al.¹⁷ obtained a maximum biomass concentration of 14.5 g/L, whereas we observed a maximum biomass concentration of 17.0 g/L in cultures of *Aurantiochytrium* sp. ATCC PRA-276 (Fig. 1(B)). This

difference is due to the substrate to biomass conversion rate ($Y_{\text{Biomass/Glucose}}$) being 0.53 in the previous study, whereas in this study, each gram of glucose consumed was converted to 0.65 g/L of biomass. The difference in the conversion rate can be attributed to the use of different species in the studies.

Cultivation using an initial TN concentration of 0.22 g/L (Fig. 1(C)) resulted in higher substrate consumption rates (0.17 g/L h glucose and 0.0022 g/L h TN) and higher cell biomass productivity (0.23 g/L h) compared with those obtained using the other TN concentrations tested in this study. In addition, the glucose to biomass conversion factor (1.28) was higher under these conditions than under the other tested conditions, which also resulted also in a higher cell concentration (23.9 g/L). Ganuza and Izquierdo¹⁸ used *Schizochytrium* sp. G13/2S to study the effect of substrate levels on lipid accumulation. These authors found that using initial concentrations of 0.30 g/L of TN and 40 g/L of glucose resulted in a biomass yield of 15.7 g/L.

In the experiments using fed-batch system (Fig. 1(D)), the concentration of TN decreased over time because its consumption rate was higher (0.0016 g/L h of TN) than its supply rate (0.0014 g/L h of TN). The concentration of glucose in the culture medium decreased until 144 h of cultivation and subsequently increased. This phenomenon can be explained by the glucose supply being greater after 144 h than that required for cell development and maintenance.

The fatty acid profiles observed in our experiments (Fig. 3) are similar to those previously obtained by Zhu et al.¹⁹ and Furlan et al.²⁰ using *Schizochytrium limacinum* OUC88 and *Thraustochytrium* sp. ATCC 26185, respectively. Zhu et al.¹⁹ found C14:0 (3.8–9.6%), C15:0 (2.1–10.1%), C16:0 (32.6–43.3%), DPA ω 6 (7.1–8.2%) and DHA (29.8–36.5%) as the main fatty acids. Furlan et al.²⁰ found C14:0 (1.5–9%), C15:0 (21–35%), C16:0 (5–33%), DPA ω 6 (7–9%) and DHA (20–31.5%) as the main fatty acids. These results are similar to our results: C14:0 (1.8–16%), C15:0 (5–16%), C16:0 (9–32.5%) and the polyunsaturated, including DPA ω 6 (6.5–14%) and DHA (20–43%). Therefore, the fatty acids produced by members of the Thraustochytriidae family are likely to be mainly C14:0, C15:0, C16:0, DPA ω 6 and DHA.

Li et al.¹⁷ studied the composition of the *Aurantiochytrium* sp. BR-MP4-A1 biomass and concluded that DPA ω 6 and DHA comprised 6.6% and 28.9%, respectively, of the total fatty acids. These authors also observed that the DPA ω 6 (18%) and DHA (78%) levels were higher than those of the other PUFAs that were quantitated, similar to the results obtained in this study (Fig. 3).

We also observed a reduction in PUFAs production relative to that of the total fatty acids when the TN concentration decreased. In contrast, C16:1 ω 9 and C18:1 ω 9 production was increased because these fatty acids are the precursors used for PUFAs synthesis. Therefore, the smaller the fractions of C16:1 ω 9 and C18:1 ω 9, the higher the fraction of PUFAs (Fig. 3).

Among the major fatty acids that are PUFAs, the content of DPA ω 6 (20–23.1%) varied little, whereas the content of DHA (61.3–70.5%) varied greatly. High concentrations of available total nitrogen in the culture medium facilitate the synthesis of other fatty acids in addition to DHA and DPA ω 6, which form a significant fraction of the PUFAs present. For example, C16:2 ω 4, C20:5 ω 3 and C22:5 ω 3 comprised 5.19%, 4.84% and

Table 1 – Total fatty acid content of the *Aurantiochytrium* sp. ATCC PRA-276 biomass with the highest PUFAs concentration under each experimental condition.

Total nitrogen	3.0 g/L	0.44 g/L	0.22 g/L	Fed-batch ^b
C/N	4	27	54	100
Culture time (h)	Total fatty acids (mg/g) ^a			
120	129.15 ± 0.15			
168		448.47 ± 0.05		
96			455.62 ± 0.09	
144				526.20 ± 0.20

^a Mean values ± standard deviation.

^b 0.14 g/L of glucose and 0.0014 g/L of total nitrogen supplied each hour.

2.06%, respectively, of the fatty acids observed in the biomass obtained in cultures grown using an initial nitrogen concentration of 3 g/L TN and C16:2 ω comprised 1.95% of the fatty acids observed in cultures grown using an initial nitrogen concentration of 0.44 g/L (Fig. 3).

The main commercial sources of PUFAs are species of fatty fish, such as herring, mackerel, salmon and sardines. The *Aurantiochytrium* strain used in this study accumulated higher concentrations of PUFAs (28–70%) than those reported in sardine oil (31.1%) by Morais.²¹ The DHA concentration (20–43%) produced by this oleaginous microorganism was 2- to 4-fold higher than that found in the sardine oil (11%).²¹ Cultivated *Aurantiochytrium* sp. ATCC PRA-276 is a promising alternative source of oil rich in PUFAs. Using fish for the large-scale production of such oil is limited by the changes in the lipid compositions and contents and the fatty acid profiles of fish, which are affected by the seasons and their species, sex, size, reproductive status, catch location, diet and nutritional status.²² Moreover, fish oil exhibits a great diversity of fatty acids with different chain lengths and degrees of unsaturation and thus, requires expensive extraction and purification processes.⁸

Cultivating an oleaginous microorganism in the laboratory under controlled environmental conditions reduces the risk of contamination and can increase fatty acid production at a low cost. In this study, we observed that the total fatty acid content of the biomass (% w/w) increased as the TN concentration was decreased. This phenomenon occurred because lipids generally accumulate in oleaginous microorganisms when the medium contains an excess of the carbon source and a limited amount of nitrogen (high C/N ratio). In the presence of low-level nitrogen, the synthesis of proteins and nucleic acids is limited by the enhanced conversion of carbon to oil.^{23,24} This process was observed in the experiments using a fed-batch system, in which C/N ratio was high, eventually leading to the accumulation of a high level of total fatty acids (526.20 mg/g) in the biomass (Table 1). However, fed-batch cultures exhibited lower PUFAs production (1.89 g/L) and consequently lower DPA ω6 (0.44 g/L) and DHA (1.33 g/L) production than those of the batch cultures because the yields of these fatty acids are dependent on the accumulation of PUFAs in the total lipids as well as on the accumulation of oils in the biomass. Additionally, the fatty acid yield was also related to the cell concentration at a given time.

For example, in the experiments using an initial TN concentration of 0.22 g/L, there was a higher substrate consumption rate, increased biomass productivity and a higher

C/N ratio (54), which resulted in higher yields of DPA ω6 (0.80 g/L) and DHA (2.54 g/L).

Ganuza and Izquierdo¹⁸ observed greater DPA ω6 (3.85% w/w) and DHA (15.4% w/w) accumulation by *Schizochytrium* sp. G13/2S cells grown using initial nitrogen and glucose concentrations of 0.30 g/L and 40 g/L, respectively, than was observed in our experiments (3.34% w/w DPA ω6, and 10.62% w/w DHA) using an initial nitrogen concentration of 0.22 g/L. However, their DPA ω6 (0.6 g/L) and DHA (2.42 g/L) production rates were lower than ours, which can be attributed to the higher maximum biomass concentration (23.9 g/L) obtained in our experiments compared with that reported by Ganuza and Izquierdo¹⁸ (15.7 g/L).

Using a similar culture medium with an initial TN concentration of 0.44 g/L to grow *Schizochytrium* sp. ATCC 20889, Jiang et al.⁸ observed that DHA accounted for 26% of the total fatty acids after 120 h of cultivation, which is very similar to the 25.5% DHA level in the total fatty acids observed in our study. In our study, the accumulated biomass consisted of approximately 12.5% (w/w) DHA, which is higher than the value (8.8%) reported by Jiang et al.⁸

Burja et al.²⁴ evaluated the effect of different concentrations of nitrogen on fatty acid production by *Thraustochytrium* sp. ONC-T18. Using an initial TN concentration of 0.75 g/L, these authors obtained 0.04 g/L of DHA, corresponding to 0.53% (w/w) of the biomass, and reported a low biomass concentration (7.5 g/L) and a low level of DHA in the biomass. Burja et al.²⁴ also observed that using a higher initial TN concentration (1.23 g/L), 1.56 g/L of DHA was obtained, which is approximately 3 times the DHA concentration (0.51 g/L DHA) observed in our experiments using an initial TN concentration of 3.0 g/L.

Conclusions

Herein, we presented the results of a study of the effect of different concentrations of carbon and nitrogen on PUFAs production by *Aurantiochytrium* sp. The PUFAs production of this microorganism depends on the accumulation of total fatty acids and on the concentration of the biomass. Therefore, the culture medium should facilitate the growth of this microorganism and provide an adequate nitrogen supply with respect to the C/N ratio because it accumulates oils when the total nitrogen supply is limited.

The main polyunsaturated fatty acids found in the *Aurantiochytrium* sp. ATCC PRA-276 biomass were DPA ω6 (20–23.1%)

and DHA (61.3–70.5%). The maximum cell concentration of 23.9 g/L (with 45.5% of its weight consisting of fatty acids) was observed at 96 h of cultivation using initial concentrations of 30 g/L of glucose and 0.22 g/L of total nitrogen. Under these conditions, the highest PUFAs concentration (3.6 g/L) was reached, with the DHA and DPA ω 6 concentrations being 2.54 and 0.80 g/L, respectively.

The results of this study showed that the growth of *Aurantiochytrium* sp. ATCC PRA-276 and its accumulation of PUFAs, particularly DHA, are dependent on the concentrations of the carbon and nitrogen substrates. The results also demonstrated that the cultivation period is an important variable for PUFAs production by *Aurantiochytrium* sp. ATCC PRA-276.

Aurantiochytrium sp. ATCC PRA-276 is capable of producing high levels of PUFAs. Therefore, developing new techniques for cultivating this microorganism could reduce the cost and increase the production of oils for use in food and medicines.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

This study was supported by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior of Brazil (CAPES) and developed at the Portuguese Institute of Sea and Atmosphere (IPMA) in Lisbon, PT, with the aid of a scholarship grant awarded to the first author by the Doctoral in the Country with Internship Abroad Programme (PDEE) (grant no. 6906/10-9). The authors also thank the ALGAENE Project and Depsiextracta Biological Technologies, Lda. for their support.

REFERENCES

1. Sijtsma L, Swaaf De ME. Biotechnological production and applications of the ω -3 polyunsaturated fatty acid docosahexaenoic acid. *Appl Microbiol Biotechnol*. 2004;64:146–153.
2. Shwu-Tzy W, Shih-Tsung Y, Liang-Ping L. Effect of culture conditions on docosahexaenoic acid production by *Schizochytrium* sp. S31. *Proc Biochem*. 2005;40:3103–3108.
3. Simopoulos AP, Kifer RR, Martin RE, Barlow SM. Health effects of omega 3 polyunsaturated fatty acids in seafoods. *World Rev Nutr Diet*. 1991;66:1–592.
4. Corsinovi L, Biasi F, Poli G, Leonarduzzi G, Isaia G. Dietary lipids and their oxidized products in Alzheimer's disease. *Mol Nutr Food Res*. 2011;55:S161–S172.
5. Nauroth JM, Liu YC, Elswyk MV, et al. Docosahexaenoic acid (DHA) and docosapentaenoic acid (DPA n-6) algal oils reduce inflammatory mediators in human peripheral mononuclear cells in vitro and paw edema in vivo. *Lipids*. 2010;45:375–384.
6. Manikan V, Kalil MS, Hamid AA. Response surface optimization of culture medium for enhanced docosahexaenoic acid production by a Malaysian thraustochytrid. *Sci Rep*. 2015;5:1–8.
7. Qu L, Ren L-J, Li J, et al. Biomass composition, lipid characterization, and metabolic profile analysis of the fed-batch fermentation process of two different docosahexaenoic acid producing *Schizochytrium* sp. strains. *Appl Biochem Biotechnol*. 2013;171:1865–1876.
8. Jiang Y, Fan KW, Wong RT, Chen F. Fatty acid composition and squalene content of the marine microalga *Schizochytrium mangrovei*. *J Agric Food Chem*. 2004;52:1196–1200.
9. Raghukumar S. Thraustochytrid marine protists: production of PUFAs and other emerging technologies. *Mar Biotechnol*. 2008;10:631–640.
10. Kendrick A, Ratledge C. Lipids of selected molds grown for production of n-3 and n-6 polyunsaturated fatty acids. *Lipids*. 1992;27:15–20.
11. Furlan VJM, Paulo MC, Batista I, Bandarra NM, Santo MLE, Prentice C. Effect of the concentration of glucose in the docosahexaenoic acid (DHA) production by *Thraustochytrium* sp. ATCC 26185. *Adv J Food Sci Technol*. 2012;4:257–264.
12. Min KH, Lee HH, Anbu P, Chaulagain BP, Hur BK. The effects of culture conditions on the growth property and docosahexaenoic acid production from *Thraustochytrium aureum* ATCC 34304. *Korean J Chem Eng*. 2012;29:1211–1215.
13. Miller GL. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal Chem*. 1959;31:426–428.
14. Cohen Z, Vonshak A, Richmond A. Effect of environmental conditions on fatty acid composition of the red algae *Porphyridium cruentum*: correlation to growth rate. *J Phycology*. 1988;24:328–332.
15. Triola MF. *Introdução à estatística*. 10th ed. Rio de Janeiro: Ltc; 2008.
16. Chen GQ, Fan KW, Lu FP. Optimization of nitrogen source for enhanced production of squalene from *Thraustochytrid Aurantiochytrium* sp. *New Biotechnol*. 2010;27:382–389.
17. Li Q, Chen GQ, Fan KW, Lu FP, Aki T, Jiang Y. Screening and characterization of squalene-producing *Thraustochytrids* from Hong Kong mangroves. *J Agric Food Chem*. 2009;57:4267–4272.
18. Ganuza E, Izquierdo MS. Lipid accumulation in *Schizochytrium* G13/2S produced in continuous culture. *Appl Microbiol Biotechnol*. 2007;76:985–990.
19. Zhu L, Zhang X, Ji L, Song X, Kuang C. Changes of lipid content and fatty acid composition of *Schizochytrium limacinum* in response to different temperatures and salinities. *Proc Biochem*. 2007;42:210–214.
20. Furlan VJM, Paulo MC, Maus VW, Ferreira J, Batista I, Bandarra NM. Production of docosahexaenoic acid (DHA) from *Thraustochytrium* sp. ATCC 26185 using different nitrogen concentrations. *B Cent Pesqui Proc A*. 2014;32:1–10.
21. Moraes MM. *Estudo do processo de refino do óleo de pescado*. Brasil: Rio Grande; 2000:100 [MSc Dissertation. Engenharia e Ciência de Alimentos, FURG].
22. Gonçalves AA. *Tecnologia do pescado: ciência, tecnologia, inovação e legislação*. São Paulo: Atheneu; 2011.
23. Ratledge C, Wynn JP. The biochemistry and molecular biology of lipid accumulation in oleaginous microorganisms. *Adv Appl Microbiol*. 2002;51:1–51.
24. Burja AM, Radianingtyas H, Windust A, Barrow CJ. Isolation and characterization of polyunsaturated fatty acid producing *Thraustochytrium* species: screening of strains and optimization of omega-3 production. *Appl Microbiol Biotechnol*. 2006;72:1161–1169.